

Compositional Evolution of Saturn's Rings: Ice, Tholin, and "Chiron"-dust

J. N. Cuzzi (Ames Research Center), P. R. Estrada (Cornell University)

We address compositional evolution in planetary ring systems subsequent to meteoroid bombardment. The huge surface area to mass ratio of planetary rings ensures the importance of this process, given currently expected values of meteoroid flux. We developed a model which includes both direct deposition of extrinsic meteoritic "pollutants", and ballistic transport of the increasingly polluted ring material as impact ejecta. Certain aspects of the observed regional variations in ring color and albedo can be understood in terms of such a process.

We conclude that the regional scale color and albedo differences between the C ring and B ring can be understood if all ring material began with the same composition (primarily water ice, based on other data, but colored by tiny amounts of non-icy, reddish absorber) and then evolved entirely by addition and mixing of extrinsic, neutrally colored, highly absorbing material. This conclusion is readily extended to the Cassini Division and its surroundings as well. Typical silicates are unable to satisfy the ring color, spectroscopic, and microwave absorption constraints either as intrinsic or extrinsic nonicy constituents. However, "Titan Tholin" provides a satisfactory match for the inferred refractive indices of the "pre-pollution" nonicy ring material. The extrinsic bombarding material is compatible with the properties of Halley or Chiron, but not with the properties of other "red" primitive objects such as Pholus.

We further demonstrate that the detailed radial profile of color across the abrupt B ring - C ring boundary is quite compatible with such a "pollution transport" process, and that the shape of the profile can constrain key parameters in the model. We use the model to estimate the "exposure age" of Saturn's rings to extrinsic meteoroid flux. We obtain a geologically young "age" which is compatible with timescales estimated independently based on the evolution of ring *structure* due to ballistic transport, and also with other "short timescales" estimated on the grounds of gravitational torques.

Abstract submitted for 1996 DPS meeting

Date submitted: LPI electronic form version 5/96

Division for Planetary Sciences Abstract Form

DPS Category 17

Running #7494

Session 0.00

Invited ☐ Poster presentation ☒ Title only ☐

Have you received your Ph.D. since the last DPS meeting?

Yes ☐ No ☐

Is your abstract newsworthy, and if so, would you be willing to prepare a news release and be available for interviews with reporters?

Yes ☐ No ☐ Maybe ☐

Paper presented by Jeffrey N. Cuzzi

Space Science Division

Ames Research Center

Mail Stop 245-3

Moffett Field CA 94035 USA

Phone: 415-604-6343

Fax: 415-604-6779

Email: cuzzi@cloud9.arc.nasa.gov

Special instructions: Tue Aug 27 17:14:35 CDT 1996

Membership Status (First Author):

DPS-AAS Member ☒ Non-Member ☐

Student Member ☐ Student Non-Member ☐

Is this your first DPS presentation? Yes ☐ No ☐

Sponsor: